

Write your name here

Surname

Other names

Centre Number

Candidate Number

**Edexcel GCE**

**Chemistry**

**Advanced Subsidiary**

**Unit 2: Application of Core Principles of Chemistry**

Friday 27 May 2011 – Afternoon

**Time: 1 hour 30 minutes**

Paper Reference

**6CH02/01**

**Candidates may use a calculator.**

Total Marks

### Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*

### Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*
- Questions labelled with an **asterisk** (\*) are ones where the quality of your written communication will be assessed  
– *you should take particular care with your spelling, punctuation and grammar, as well as the clarity of expression, on these questions.*
- A Periodic Table is printed on the back cover of this paper.

### Advice

- Read each question carefully before you start to answer it.
- Keep an eye on the time.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P38479A

©2011 Edexcel Limited.

7/7/5/5/3/







P 3 8 4 7 9 A 0 1 2 0

**edexcel**  
advancing learning, changing lives

## SECTION A

Answer ALL the questions in this section. You should aim to spend no more than 20 minutes on this section. For each question, select one answer from A to D and put a cross in the box . If you change your mind, put a line through the box  and then mark your new answer with a cross .

1 The correct balanced equation for the reaction between heated magnesium and steam, including state symbols, is





(Total for Question 1 = 1 mark)

2 This question concerns the trends in properties on descending Group 2 of the Periodic Table.



(a) What are the trends in solubility of sulfates and hydroxides down Group 2?

(1)





(b) What are the trends in thermal stability of carbonates and nitrates down Group 2?

(1)



(c) What are the trends in first ionization energy and electronegativity of the elements down Group 2?

(1)



(Total for Question 2 = 3 marks)



3 Which silver halide is a cream coloured solid which darkens in sunlight and dissolves in concentrated ammonia solution?

- A AgF
- B AgCl
- C AgBr
- D AgI

(Total for Question 3 = 1 mark)

4 What is the FBF bond angle in boron trifluoride,  $\text{BF}_3$ ?

- A  $180^\circ$
- B  $120^\circ$
- C  $109.5^\circ$
- D  $90^\circ$

(Total for Question 4 = 1 mark)

5 What is the total number of electrons in the covalent bonds in a beryllium chloride molecule,  $\text{BeCl}_2$ ?

- A 2
- B 4
- C 6
- D 8

(Total for Question 5 = 1 mark)

6 Which of the following molecules is linear?

- A  $\text{CO}_2$
- B  $\text{C}_2\text{H}_4$
- C  $\text{H}_2\text{O}$
- D  $\text{NH}_3$

(Total for Question 6 = 1 mark)



7 Which of the following molecules is **non-polar**?

- A  $\text{CH}_3\text{Cl}$
- B  $\text{CH}_2\text{Cl}_2$
- C  $\text{CHCl}_3$
- D  $\text{CCl}_4$

(Total for Question 7 = 1 mark)

8 Methanol dissolves in water mainly due to the formation of new

- A hydrogen bonds.
- B dipole-dipole forces.
- C London forces.
- D covalent bonds.

(Total for Question 8 = 1 mark)

9 Which of the following molecules does **not** absorb infrared radiation?

- A  $\text{N}_2$
- B  $\text{NO}_2$
- C  $\text{CO}$
- D  $\text{CO}_2$

(Total for Question 9 = 1 mark)

10 There would be a major peak in the mass spectrum for butan-1-ol,  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$ , but **not** for butan-2-ol,  $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$ , at  $m/e$  value

- A 15
- B 17
- C 29
- D 43

(Total for Question 10 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



11 How many molecular ion peaks (parent ion peaks) occur in the mass spectrum of 1,2-dibromoethane,  $\text{CH}_2\text{BrCH}_2\text{Br}$ ?

Assume the only isotopes present are  $^1\text{H}$ ,  $^{12}\text{C}$ ,  $^{79}\text{Br}$  and  $^{81}\text{Br}$ .

- A 1
- B 2
- C 3
- D 4

(Total for Question 11 = 1 mark)

12 The following reactions have been used in the chemical industry to make liquid and solid products, allowing any gaseous products to escape into the atmosphere:

- A  $\text{CH}_3\text{OH}(\text{g}) + \text{CO}(\text{g}) \rightarrow \text{CH}_3\text{COOH}(\text{l})$
- B  $\text{CaCO}_3(\text{s}) \rightarrow \text{CaO}(\text{s}) + \text{CO}_2(\text{g})$
- C  $\text{CH}_4(\text{g}) + 3\text{Cl}_2(\text{g}) \rightarrow \text{CHCl}_3(\text{l}) + 3\text{HCl}(\text{g})$
- D  $\text{CH}_2\text{CH}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightarrow \text{CH}_2\text{ClCH}_2\text{Cl}(\text{l})$

(a) Which reaction has an atom economy by mass of 56%?

(1)

- A
- B
- C
- D

(b) Which reaction causes the most immediate damage to the environment?

(1)

- A
- B
- C
- D

(c) Which reaction is an electrophilic addition?

(1)

- A
- B
- C
- D

(Total for Question 12 = 3 marks)



13 Propan-1-ol and propan-2-ol are separately oxidized under mild conditions by acidified sodium dichromate(VI) and the product immediately distilled off. What is the oxidation product in each case?

		Propan-1-ol	Propan-2-ol
<input checked="" type="checkbox"/>	A	propanal	propanone
<input checked="" type="checkbox"/>	B	propanoic acid	propanone
<input checked="" type="checkbox"/>	C	propanal	propanoic acid
<input checked="" type="checkbox"/>	D	propanone	propanal

(Total for Question 13 = 1 mark)

14 Unsaturated vegetable oils are hardened to make margarine by reaction with hydrogen and a nickel catalyst. Which terms could both be used to describe this type of reaction?

- A Substitution and oxidation
- B Substitution and reduction
- C Addition and oxidation
- D Addition and reduction

(Total for Question 14 = 1 mark)

15 When iodomethane,  $\text{CH}_3\text{I}$ , is heated in a sealed tube with an excess of alcoholic ammonia, which of the following **cannot** be formed?

- A Methylamine,  $\text{CH}_3\text{NH}_2$
- B Ethylamine,  $\text{CH}_3\text{CH}_2\text{NH}_2$
- C Dimethylamine,  $(\text{CH}_3)_2\text{NH}$
- D Ammonium iodide,  $\text{NH}_4\text{I}$

(Total for Question 15 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



- 16 The enthalpy change of neutralization of an acid by an alkali is measured by adding  $10.0 \text{ cm}^3$  of hydrochloric acid to  $10.0 \text{ cm}^3$  of sodium hydroxide.  $10.0 \text{ cm}^3$  pipettes with an accuracy of  $\pm 0.04 \text{ cm}^3$  are used to measure out both solutions.

The overall percentage error in measuring the total volume of the reaction mixture is

- A  $\pm 0.04\%$
- B  $\pm 0.08\%$
- C  $\pm 0.4\%$
- D  $\pm 4.0\%$

(Total for Question 16 = 1 mark)

---

**TOTAL FOR SECTION A = 20 MARKS**



D 3 8 4 7 0 A 0 7 2 0

## SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

17 This question is about the element chlorine and its compounds.

(a) When chlorine is bubbled through water, a solution of chlorine water forms. What is the colour of chlorine water?

(1)

(b) Chlorine water is added to potassium iodide solution.

(i) State the colour of the solution produced.

(1)

(ii) Write the **ionic** equation for the reaction, including state symbols.

(2)

(c) The concentration of chlorine water was found by taking  $10.0 \text{ cm}^3$  of solution, adding an excess of potassium iodide solution, and titrating with  $0.0100 \text{ mol dm}^{-3}$  of sodium thiosulfate solution. The experiment was repeated.

The following results were obtained.

Titration number	1	2
Final burette reading / $\text{cm}^3$	38.60	47.60
Initial burette reading / $\text{cm}^3$	29.50	38.60
Volume added / $\text{cm}^3$	9.10	9.00





- (i) Name a suitable indicator for the titration. State the colour change you would expect to see at the end point.

(2)

Indicator .....

Colour change from ..... to .....

- (ii) Calculate the mean titre and use this value to calculate the number of moles of sodium thiosulfate used in the titration.


(1)

Mean titre = .....  $\text{cm}^3$

Moles of sodium thiosulfate

- (iii) Complete the ionic equation for the reaction between iodine and thiosulfate ions.

(2)



- (iv) Calculate the number of moles of iodine which reacted with the sodium thiosulfate solution.

(1)

- (v) Hence state the number of moles of chlorine present in  $10.0 \text{ cm}^3$  of the chlorine water.

(1)

- (vi) Calculate the concentration of the chlorine water, in  $\text{mol dm}^{-3}$ .

(1)



(d) Potassium burns in chlorine to form potassium chloride.

(i) Give the colour of the flame when potassium burns in chlorine.

(1)

(ii) Write the equation for the reaction between potassium and chlorine. State symbols are **not** required.

(1)

(e) Concentrated sulfuric acid is added to potassium chloride in a test tube. Steamy fumes are given off which react with ammonia to give dense white smoke.

(i) Name the gas given off in this reaction.

(1)

(ii) Steamy fumes are observed at the mouth of the test tube. Explain how these fumes are formed.

(1)

(iii) The steamy fumes react with ammonia to give a dense white smoke. Identify the white smoke by name or formula.

(1)

(f) 2-chlorobutane can be made from butan-2-ol.

(i) Name the chemical you would add to butan-2-ol in the laboratory to make 2-chlorobutane.

(1)



- (ii) 2-chlorobutane reacts with alcoholic potassium hydroxide at a high temperature to form a mixture of gaseous alkenes.

Draw a fully labelled diagram of the apparatus you would use to prepare and collect this mixture.

(3)

---

(Total for Question 17 = 21 marks)



**BLANK PAGE**



18 This question is about ethanethiol,  $\text{CH}_3\text{CH}_2\text{SH}$ . Thiols are like alcohols, but the oxygen atom has been replaced by a sulfur atom. They react in a similar way to alcohols.

(a) (i) Draw a dot and cross diagram for ethanethiol, showing outer electrons only. (2)

(ii) Give the value for the CSH bond angle in ethanethiol. Justify your answer. (3)

CSH angle .....

Justification .....

.....

.....

.....

(b) There are hydrogen bonds between ethanol molecules but not between ethanethiol molecules.

(i) Explain why the bond angle around the hydrogen atom involved in a hydrogen bond is  $180^\circ$ . (2)

.....

.....

.....

(ii) Explain why there are no hydrogen bonds between ethanethiol molecules. (1)

.....

.....



(c) (i) Describe the formation of London forces.

(2)

(ii) Explain why the London forces in ethanethiol are stronger than those in ethanol.

(1)

(d) The reaction of sodium with ethanethiol,  $\text{CH}_3\text{CH}_2\text{SH}$ , is similar to its reaction with ethanol.

(i) Suggest one observation you would make when sodium is added to ethanethiol.

(1)

(ii) Suggest a balanced equation for this reaction. State symbols are **not** required.

(1)



(e) Ethanol can be made from bromoethane by reaction with aqueous potassium hydroxide, KOH(aq), under suitable conditions.

(i) Write the equation for this reaction. State symbols are **not** required.

(1)

(ii) State the type and mechanism of this reaction.

(2)

Type .....

Mechanism .....

(iii) Suggest the formula of a suitable chemical to make ethanethiol from bromoethane.

(1)

(f) When ethanethiol undergoes complete combustion in air, a gas is produced which is not formed on the complete combustion of ethanol. Identify the gas and suggest why it is damaging to the environment.

(2)

(Total for Question 18 = 19 marks)

**TOTAL FOR SECTION B = 40 MARKS**



## SECTION C

Answer ALL the questions. Write your answers in the spaces provided.

- 19 This question is about nitrogen monoxide, NO, which can be described both as a friend and a foe.

Chemists have discovered that nitrogen monoxide plays an important role in the body by dilating blood vessels. If someone is suffering from blood circulatory or heart problems, a chemical may be given which will quickly break down to give nitrogen monoxide. Years ago, nitroglycerine was used for this purpose. Interestingly, the same chemical Nobel had used to make dynamite was used to treat him in old age.

In the laboratory, nitrogen monoxide can be prepared by adding concentrated nitric acid to powdered silver. Nitrogen monoxide is a colourless gas which is partially soluble in water. It is difficult to detect its smell, because it reacts with oxygen in the air to form pungent-smelling nitrogen dioxide.

Nitrogen monoxide is formed when a mixture of air and oxygen is heated to a high temperature. This reaction occurs in the engines of cars and aeroplanes. Nitrogen monoxide has a disastrous effect on the ozone layer because it is a free radical. Nitrogen monoxide is also a greenhouse gas.

- (a) (i) What is meant by the term **free radical**?

(1)


- (ii) Suggest a dot and cross diagram for nitrogen monoxide, showing outer shell electrons only, remembering that it is a free radical.

(2)





- (b) (i) Part of the **unbalanced** equation for the preparation of nitrogen monoxide from nitric acid is shown below.



Identify the elements which are oxidized and reduced and give their oxidation numbers.

(3)

Element oxidized .....


Oxidation number initial ..... final .....

Element reduced .....

Oxidation number initial ..... final .....

- (ii) Complete and balance the equation for the reaction between silver and nitric acid.

(2)



- (c) The reaction between nitrogen and oxygen to form nitrogen monoxide reaches equilibrium.



- (i) Explain why the yield of nitrogen monoxide is increased when the temperature is increased.

(1)

- .....  
.....  
\*(ii) State and explain the effect, if any, on the yield of nitrogen monoxide when the pressure is increased.

(2)



(ii) State and explain how the rate of the reaction is affected by an increase in pressure.

(2)

.....

.....

.....

.....

.....

.....

.....

.....

\*(d) (i) Explain why a jet aeroplane in flight causes much more damage to the ozone layer than cars carrying the same number of passengers at sea level. You should assume that the nitrogen monoxide outputs for both methods of conveying the passengers are the same.

(2)

.....



.....

.....

.....



(ii) The reactions of chlorine free radicals with ozone may be represented by the following equations.



Write corresponding equations for the reactions of the free radical nitrogen monoxide with ozone. Combine your two equations to show the overall reaction.

Use these equations to explain why a small quantity of nitrogen monoxide can have a continuing effect on the ozone layer.

(5)

Equations

Explanation .....

---

(Total for Question 19 = 20 marks)

---

**TOTAL FOR SECTION C = 20 MARKS**  
**TOTAL FOR PAPER = 80 MARKS**



# The Periodic Table of Elements

	1	2	3	4	5	6	7	0 (8)																																																				
	6.9 <b>Li</b> lithium 3	9.0 <b>Be</b> beryllium 4	<div style="border: 1px solid black; padding: 5px; margin: 5px auto; width: fit-content;">                     1.0 <b>H</b> hydrogen 1                 </div>						4.0 <b>He</b> helium 2																																																			
	23.0 <b>Na</b> sodium 11	24.3 <b>Mg</b> magnesium 12	<div style="border: 1px solid black; padding: 5px; margin: 5px auto; width: fit-content;">                     relative atomic mass atomic symbol name atomic (proton) number                 </div>						20.2 <b>Ne</b> neon 10																																																			
(1)	39.1 <b>K</b> potassium 19	40.1 <b>Ca</b> calcium 20	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)																																										
	85.5 <b>Rb</b> rubidium 37	87.6 <b>Sr</b> strontium 38	45.0 <b>Sc</b> scandium 21	47.9 <b>Ti</b> titanium 22	50.9 <b>V</b> vanadium 23	52.0 <b>Cr</b> chromium 24	54.9 <b>Mn</b> manganese 25	55.8 <b>Fe</b> iron 26	58.9 <b>Co</b> cobalt 27	58.7 <b>Ni</b> nickel 28	63.5 <b>Cu</b> copper 29	65.4 <b>Zn</b> zinc 30	10.8 <b>B</b> boron 5	12.0 <b>C</b> carbon 6	14.0 <b>N</b> nitrogen 7	16.0 <b>O</b> oxygen 8	19.0 <b>F</b> fluorine 9	4.0 <b>He</b> helium 2																																										
	132.9 <b>Cs</b> caesium 55	137.3 <b>Ba</b> barium 56	138.9 <b>La*</b> lanthanum 57	178.5 <b>Hf</b> hafnium 72	180.9 <b>Ta</b> tantalum 73	183.8 <b>W</b> tungsten 74	186.2 <b>Re</b> rhenium 75	190.2 <b>Os</b> osmium 76	192.2 <b>Ir</b> iridium 77	195.1 <b>Pt</b> platinum 78	197.0 <b>Au</b> gold 79	200.6 <b>Hg</b> mercury 80	27.0 <b>Al</b> aluminium 13	28.1 <b>Si</b> silicon 14	31.0 <b>P</b> phosphorus 15	32.1 <b>S</b> sulfur 16	35.5 <b>Cl</b> chlorine 17	39.9 <b>Ar</b> argon 18																																										
	[223] <b>Fr</b> francium 87	[226] <b>Ra</b> radium 88	[227] <b>Ac*</b> actinium 89	[261] <b>Rf</b> rutherfordium 104	[262] <b>Db</b> dubnium 105	[266] <b>Sg</b> seaborgium 106	[264] <b>Bh</b> bohrium 107	[277] <b>Hs</b> hassium 108	[268] <b>Mt</b> meitnerium 109	[271] <b>Ds</b> darmstadtium 110	[272] <b>Rg</b> roentgenium 111	114.8 <b>In</b> indium 49	118.7 <b>Sn</b> tin 50	121.8 <b>Sb</b> antimony 51	127.6 <b>Te</b> tellurium 52	126.9 <b>I</b> iodine 53	[210] <b>At</b> astatine 85	[222] <b>Rn</b> radon 86																																										
	* Lanthanide series * Actinide series												Elements with atomic numbers 112-116 have been reported but not fully authenticated																																															
	140 <b>Ce</b> cerium 58	141 <b>Pr</b> praseodymium 59	144 <b>Nd</b> neodymium 60	150 <b>Sm</b> samarium 62	152 <b>Eu</b> europium 63	157 <b>Gd</b> gadolinium 64	159 <b>Tb</b> terbium 65	163 <b>Dy</b> dysprosium 66	165 <b>Ho</b> holmium 67	167 <b>Er</b> erbium 68	169 <b>Tm</b> thulium 69	173 <b>Yb</b> ytterbium 70	175 <b>Lu</b> lutetium 71	[254] <b>Fm</b> fermium 100	[255] <b>Es</b> einsteinium 99	[256] <b>Md</b> mendelevium 101	[257] <b>Lr</b> lawrencium 103	[258] <b>No</b> nobelium 102	[259] <b>U</b> uranium 92	[260] <b>Np</b> neptunium 93	[261] <b>Pu</b> plutonium 94	[262] <b>Am</b> americium 95	[263] <b>Cm</b> curium 96	[264] <b>Bk</b> berkelium 97	[265] <b>Cf</b> californium 98	[266] <b>Bk</b> berkelium 97	[267] <b>Cm</b> curium 96	[268] <b>Bk</b> berkelium 97	[269] <b>Cf</b> californium 98	[270] <b>Bk</b> berkelium 97	[271] <b>Cm</b> curium 96	[272] <b>Bk</b> berkelium 97	[273] <b>Cf</b> californium 98	[274] <b>Bk</b> berkelium 97	[275] <b>Cf</b> californium 98	[276] <b>Bk</b> berkelium 97	[277] <b>Cf</b> californium 98	[278] <b>Bk</b> berkelium 97	[279] <b>Cf</b> californium 98	[280] <b>Bk</b> berkelium 97	[281] <b>Cf</b> californium 98	[282] <b>Bk</b> berkelium 97	[283] <b>Cf</b> californium 98	[284] <b>Bk</b> berkelium 97	[285] <b>Cf</b> californium 98	[286] <b>Bk</b> berkelium 97	[287] <b>Cf</b> californium 98	[288] <b>Bk</b> berkelium 97	[289] <b>Cf</b> californium 98	[290] <b>Bk</b> berkelium 97	[291] <b>Cf</b> californium 98	[292] <b>Bk</b> berkelium 97	[293] <b>Cf</b> californium 98	[294] <b>Bk</b> berkelium 97	[295] <b>Cf</b> californium 98	[296] <b>Bk</b> berkelium 97	[297] <b>Cf</b> californium 98	[298] <b>Bk</b> berkelium 97	[299] <b>Cf</b> californium 98	[300] <b>Bk</b> berkelium 97

